



NPS55-78-022

NAVAL POSTGRADUATE SCHOOL

Monterey, California





DISCRETE TIME SERIES GENERATED

BY MIXTURES III:

AUTOREGRESSIVE PROCESSES (DAR(p))

IVE PROCESSES (DAR (p)

by

P. A. Jacobs

and

P. A. W. Lewis

September 1978

Approved for public release; distribution unlimited.

78 12 11 172

NAVAL POSTGRADUATE SCHOOL MONTEREY, CALIFORNIA

Rear Admiral T. F. Dedman Superintendent

J. R. Borsting Provost

Reproduction of all or part of this report is authorized.

This report was prepared by:

P. A. Jacobs, Associate Professor Department of operations Research

P. A. W. Lewis, Professor

Department of Operations Research and Statistics

Reviewed by:

Released by:

Michael G. Sovereign, Chairman

Department of Operations Research

William M. Tolles

Dean of Research

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION	READ INSTRUCTIONS BEFORE COMPLETING FORM										
REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER									
NPS55-78-022											
4. TITLE (and Subtitio)	1	TYPE OF REPORT & PERIOD COVERED									
Discrete Time Series Generated by	y Mixtures, III.	9 Technical rept.									
Autoregressive Processes (DAR(p)) .	U									
+ + A039467		6. PERFORMING ORG. REPORT NUMBER									
1 1001767											
7. AUTHOR(*)		8. CONTRACT OR GRANT NUMBER(#)									
P. A. Jacobs and P. A. W. Lewis											
9. PERFORMING ORGANIZATION NAME AND APPRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS									
Naval Postgraduate School											
Monterey, California 93940		61153N , RR014-05-01									
		N 0001478WR80012									
11. CONTROLLING OFFICE NAME AND ADDRESS		12 REPORT DATE									
Office of Naval Research		September 1978									
Arlington, Virginia 22217		TS. NUMBER OF PAGES									
14. MONITORING AGENCY NAME & ADDRESS(If differen	t from Controlling Office)	15. SECURITY CLASS. (of this report)									
	-										
N6)RRØ14Ø5 (17)RA	10740507	Unclassified									
Lancas de la companya del companya del la companya del companya de la companya del companya de la companya del la companya del la companya del la companya de la companya de la companya de la companya de la companya del l	and the same of th	15a. DECLASSIFICATION/DOWNGRADING									
		SCHEDULE									
16. DISTRIBUTION STATEMENT (of this Report)											
Approved for public release; dist	tribution unlimi	ted.									
(12) 41 p. (15) NSF-47-898285 NSF-AF-476											
17. DISTRIBUTION STATEMENT (of the abstract entered	in Block 20, If allierent tro	in Report)									
18. SUPPLEMENTARY NOTES											
19. KEY WORDS (Continue on reverse side if necessary an											
Discrete Time Series	Discrete Ran										
Mixtures	Marginal Dis										
Autoregressive Processes	Correlation :										
Markov Chain	DAR(p) Proces	SS									
Stationary Sequence											
	d identify by block number										
A scheme for obtaining a station	d identify by block number) ary sequence of	discrete random variables									
A scheme for obtaining a station	ary sequence of	discrete random variables									
A scheme for obtaining a stational which has p-th order Markov dependent	ary sequence of endence and a spe	discrete random variables ecified marginal distribution									
A scheme for obtaining a stational which has p-th order Markov depends presented. This DAR(p) process	ary sequence of e endence and a spe ss, which is a pa	discrete random variables ecified marginal distribution articular p-th order Markov									
A scheme for obtaining a stational which has p-th order Markov dependence of the presented. This DAR(p) process chain, has the physical and correspond to the physical and the physica	ary sequence of endence and a sposs, which is a posterior structure	discrete random variables ecified marginal distribution articular p-th order Markov e of an autoregressive process									
A scheme for obtaining a stational which has p-th order Markov deposis presented. This DAR(p) process chain, has the physical and corresponded of order p. The process and its	ary sequence of endence and a spense, which is a period and the end of the en	discrete random variables ecified marginal distribution articular p-th order Markov e of an autoregressive process ix are determined by the									
A scheme for obtaining a stational which has p-th order Markov dependence of the presented. This DAR(p) process chain, has the physical and correspond to the physical and the physica	ary sequence of endence and a spense, which is a period attention structure transition matriand by several or	discrete random variables ecified marginal distribution articular p-th order Markov e of an autoregressive process ix are determined by the ther parameters which, inde-									

DD 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601 SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

251 450

ECURITY CLASSIFICATION OF THIS PAGE(When Date Entered) 20 Continued. Correlational properties and initial conditions for stationarity of the process are studied. Asymptotic properties for the process are also obtained, ACCESSION for

NTIS NAME Caction DOC But Section D UNANHOUSEDED BY BISTRIBUTION/AVAILABILITY CODES Dist. AVAIL. and/or SPECIAL

DISCRETE TIME SERIES GENERATED BY MIXTURES III: AUTOREGRESSIVE PROCESSES (DAR(p))

by

P. A. Jacobs*

and

P. A. W. Lewis

Naval Postgraduate School Monterey, California

0. ABSTRACT

A scheme for obtaining a stationary sequence of discrete random variables which has p-th order Markov dependence and a specified marginal distribution is presented. This DAR(p) process, which is a particular p-th order Markov chain, has the physical and correlation structure of an autoregressive process of order p. The process and its transition matrix are determined by the specified marginal distribution and by several other parameters which, independently of the marginal distribution, determine the correlation structure. Correlational properties and initial conditions for stationarity of the process are studied. Asymptotic properties for the process are also obtained.

Support under National Science Foundation Grant NSF-77-09020 is gratefully acknowledged.

^{**} Support under Office of Naval Research Grant NR \$\text{\theta}42-284\$ and National Science Foundation Grant AF476 is gratefully acknowledged.

1. INTRODUCTION

In Jacobs and Lewis (1978a) a scheme for obtaining a sequence of discrete random variables with first order Markov dependence and with a given marginal distribution is presented. The process is a Markov chain and has the physical and correlation structure of a first order autoregressive process. In this paper we extend this result and present a scheme for obtaining a stationary sequence of discrete random variables with p-th order Markov dependence and having a given marginal distribution. The process is a p-th order Markov chain and has the physical and correlation structure of a higher order autoregressive process, i.e. X_n depends explicitly on X_{n-1}, \ldots, X_{n-p} . As with the models in Jacobs and Lewis (1978a) the process is specified by the stationary marginal distribution of X_n and several other chosen parameters which, independently of the marginal distribution, determine the correlation structure.

This DAR(p) process is defined as follows. Let $\{Y_n\}$ be a sequence of independent identically distributed random variables taking values in E a discrete subset of the real line with distribution π . Let $\{V_n\}$ be a sequence of independent $\{0,1\}$ -random variables with

$$P\{v_n = 1\} = 1 - P\{v_n = 0\} = \rho$$

for $0 \le \rho < 1$. Let $\{A_n\}$ be a sequence of independent random variables taking values in $\{1,2,\ldots,p\}$ with $P\{A_n=i\}=\alpha_i \ge 0$, $i=1,2,\ldots,p$ where $\sum_{i=1}^p \alpha_i=1$. Put

(1.1)
$$X_n = V_n X_{n-A_n} + (1 - V_n) Y_n$$

for $n=1,2,\ldots$. Then $\{X_n\}$ is called the DAR(p) process (discrete autoregressive process of order p). It will be shown in Section 3 that it is possible to obtain an initial distribution for $(X_{-p+1},\ldots,X_1,X_0)$ which makes the sequence $\{X_n\}$ stationary with marginal distribution π .

In Section 2 we will study the correlational structure of the stationary DAR(p) process, obtaining Yule-Walker type equations and discussing their solution for the DAR(2) and DAR(3) process.

Section 3 discusses the asymptotic properties of the DAR(p) process; it is shown that the process is ϕ -mixing in the sense of Billingsley (1968) and consequently that, with the usual definitions, sample means, covariances and quantiles converge almost surely and are asymptotically normal.

The process as defined at (1.1) has the property that all of the serial correlations $r(k) = corr(X_{n+k}, X_n)$, $k = \pm 1, \pm 2, \ldots$, are greater than or equal to zero. This problem can be overcome quite simply if π is a symmetric distribution. The process then becomes a full analog to the standard linear AR(p) process (see e.g., Box and Jenkins, 1970). A discussion of the correlation properties of this extended DAR(p) process is given. When π is not symmetric it is necessary to consider schemes such as coupling two DAP(p) processes over a bivariate sequence $\{Y_n, Y_n^*\}$, where

the Y_n 's and Y_n 's are negatively correlated. Since this leads into the question of multivariate DAR(p) processes, it will be explicated elsewhere.

2. CORRELATIONAL PROPERTIES

Let $\{X_n\}$ be a stationary DAR(p) process having marginal distribution π and parameters ρ , α_i , $i=1,\ldots,p$, and let $m=E(Y_n)=E(X_n)$. From (1.1) it follows that the dependence in the DAR(p) process is due to the random selection procedure by which X_n chooses one of the Y_k ; $k \leq n$. In other words it is possible to write X_n as a random index model over Y_k , $k \leq n$, although the indexing variables for each n are correlated. To derive the correlation structure it is easier to work with the autoregressive formulation (1.1). Note that with probability $\rho\alpha_j$, $X_{n+k} = X_{n+k-j}$ for $j=1,\ldots,p$. This observation allows one to derive Yule-Walker equations for the serial correlations $r(k) = Corr(X_{n+k}, X_n)$ quite simply. For example, if we center the X_n 's to give $X_n' = X_n - m$, then multiplying (1.1) by X_{n-1}' and taking expectations

$$\begin{split} r(1) &= E(X_{n}^{\dagger}X_{n-1}^{\dagger}) \\ &= E(X_{n-1}^{\dagger}V_{n}X_{n-A_{n}}^{\dagger}) + E(X_{n-1}^{\dagger}(1-V_{n})Y_{n}^{-m}) \\ &= \rho\alpha_{1}r(0) + \rho\alpha_{2}r(1) + \cdots + \rho\alpha_{p}r(p-1). \end{split}$$

Continuing this way, the complete set of Yule-Walker equations is obtained:

(2.1)
$$r(1) = \rho \alpha_1 r(0) + \rho \alpha_2 r(1) + \cdots + \rho \alpha_p r(p-1)$$
,

(2.2)
$$r(2) = \rho \alpha_1 r(1) + \rho \alpha_2 r(0) + \cdots + \rho \alpha_p r(p-2)$$
,
: : : :

(2.3)
$$r(p) = \rho \alpha_1 r(p-1) + \rho \alpha_2 r(p-2) + \cdots + \rho \alpha_p r(0)$$
,

and, for k > 1,

(2.4)
$$r(p+k) = \rho \alpha_1 r(p+k-1) + \rho \alpha_2 r(p+k-2) + \cdots + \rho \alpha_p r(k)$$
.

Clearly, when ρ = 0 all r(k)'s, k = 1,2,... are zero, so that one is interested in solving the equations for 0 < ρ < 1.

To see that the serial correlations are all positive or zero, let q(i) be the probability that X_n and X_{n+1} choose the same random variable Y_k , where, because of the backward definition of the autoregression, $k \leq n$. Then q(i) also satisfies equations (2.1)-(2.4). Hence r(i)=q(i), $i=1,2,\ldots$ and the correlations for the DAR(p) process are always nonnegative.

If we define a generating function

$$\beta(z) = \sum_{j=1}^{\infty} z^{j} r(j)$$
 for $0 \le z$,

then from (2.1) - (2.4) we have that

(2.5)
$$\beta(z)$$

$$= \{z[\rho\alpha_{1}r(0) + \cdots + \rho\alpha_{p}r(p-1)] + z^{2}[\rho\alpha_{2}r(0) + \cdots + \rho\alpha_{p}r(p-2)] + z^{p}[\rho\alpha_{p}r(0)]\} \{1 - z\rho\alpha_{1} - z^{2}\rho\alpha_{2} - \cdots - z^{p}\rho\alpha_{p}\}^{-1}.$$

The roots of the characteristic equation

(2.6)
$$1 - z_{\rho\alpha_1} - z_{\rho\alpha_2}^2 - \cdots - z_{\rho\alpha_p}^p = 0$$

can be used to expand $\beta(z)$ by partial fractions to obtain expressions for r(k). The roots can also be used to characterize the behavior of the autoregressive function $\{r(k)\}$. This will be illustrated for the DAR(2) and DAR(3) processes.

First we note that since ρ and the α_i 's are greater than or equal to zero, the left-hand side of (2.6) equals 1 at z=0, $(1-\rho)>0$ at z=1, and since it is monotone in z for $z\geq 0$, there is one and only one positive root for a value of z>1. All other roots are negative or imaginary. (For p=1, the root is $z_1=1/\rho$.)

Another consequence of (2.6) concerns the quantity c defined in Jacobs and Lewis (1978b). We have

$$\lim_{m \to \infty} m \operatorname{var}\left(\frac{X_1 + \cdots + X_m}{m}\right) = \operatorname{var}(X_1) c,$$

where

$$c = 1 + 2 \sum_{j=1}^{\infty} r(j)$$
.

From (2.5)

(2.7)
$$c = 1 + 2\beta(1)$$

$$= 1 + 2\rho\{r(0) + (1-\alpha_1) \quad r(1) + (1-\alpha_1-\alpha_2) \quad r(2)$$

$$+ \cdots + (1-\alpha_1 - \cdots - \alpha_{p-1}) \quad r(p-1)\}/(1-\rho)$$

$$\geq 1 + \frac{2\rho}{1-\rho}.$$

The inequality follows because all of the serial correlations r(k) are nonnegative.

Example 2.1. The DAR(2) process.

In the definition (1.1) let $P\{A_n=1\}=1-P\{A_n=2\}=\alpha$ for $0\leq\alpha\leq1$. If $\alpha=1$ the DAR(2) process reduces to the DAR(1) process which was studied in Jacobs and Lewis (1978 a,b). The Yule-Walker equations are easily solved to give

(2.7)
$$r(1) = \rho \alpha [1 - \rho (1-\alpha)]^{-1}$$

and

(2.8)
$$\mathbf{r}(2) = \{\rho(1-\rho) + 2\rho\alpha[\rho-1]\}\{1 - \rho(1-\alpha)\}^{-1}$$

$$= \mathbf{r}(1)[(1-\rho)\alpha^{-1} + 2\rho - 1] .$$

If $\alpha=1$, $r(1)=\rho$, $r(2)=\rho^2$, ... as it should for a first order Markov chain; if $\alpha=0$, then r(1)=0, $r(2)=\rho$, r(3)=0, $r(4)=\rho^2$, To obtain possible values of r(1) and r(2) note that from (2.7)

$$r(1) (1-r(1))^{-1} = \alpha \rho (1-\rho)^{-1}$$
.

Thus for fixed r(1), ρ must be greater than or equal to r(1) since $0 < \alpha \le 1$. Further, for fixed r(1) and $\rho \ge r(1)$

(2.9)
$$\alpha = r(1) [1 - r(1)]^{-1} (1-\rho) \rho^{-1}$$
.

Suppose r(1) is fixed and $\rho \geq r(1)$; then substituting (2.9) into (2.8) yields

$$r(2) = -r(1) + \rho[1 + r(1)] \ge r(1)^2$$
.

It now follows that the admissible region for r(1) and r(2) is $1 > r(2) > r(1)^2$.

Note that

(2.10)
$$\frac{r(1) + r(2)}{1 + r(1)} = \rho$$

and

$$\rho\alpha = \frac{r(1) [1 + r(2)]}{1 - r(1)^2}.$$

Hence, it is possible to obtain estimates for $\,\rho\,$ and $\,\alpha\,$ from estimates of the first two serial correlations.

To study the autoregressive function note that the characteristic equation (2.6) becomes

$$1 - z\rho\alpha - z^2\rho(1-\alpha) = 0$$

which has two real roots, for $0 \le \alpha \le 1$,

$$z_{1,2} = \{\rho\alpha + a\}\{-2\rho(1-\alpha)\}^{-1},$$

where

$$a = [(\rho \alpha)^2 + 4\rho(1-\alpha)]^{1/2}$$
.

Since a $\geq [(\rho\alpha)^2]^{1/2} = \rho\alpha$, one root is positive, and the other is negative. Also the negative root is dominant in absolute value. Thus, the autocorrelation function for the DAR(2) process is nonnegative and is a mixture of the larger inverse of the positive root decaying geometrically and the smaller (in absolute value) inverse of the negative root decaying in absolute value geometrically but oscillating in sign. Specifically,

(2.12)
$$r(k) = b_1 z_1^{-k} + b_2 z_2^{-k}$$

for b_1 and b_2 given as

$$b_1 = z_1^{-1}[1 - z_2^{-2}] \{(z_1^{-1} - z_2^{-1})(1 + z_1^{-2}z_2^{-2})\}^{-1}$$

$$b_2 = z_2^{-1}(1 - z_1^{-2}) \{(z_1^{-1} - z_2^{-1})(1 + z_1^{-1}z_2^{-2})\}^{-1}.$$

Eventually the larger inverse root z_2^{-1} dominates in r(k) and the decay is geometric.

Example 2.2. The DAR(3) process.

In definition (1.1) let

$$P\{A_n=1\} = \alpha_1, \qquad P\{A_n=2\} = \alpha_2, \qquad P\{A_n=3\} = 1 - \alpha_1 - \alpha_2$$

for $0 \le \alpha_1$, $0 \le \alpha_2$, $0 \le \alpha_1 + \alpha_2 \le 1$. If $\alpha_1 + \alpha_2 = 1$ then the process reduces to the DAR(2) process. The Yule-Walker equations become

(2.13)
$$r(1) = \rho \alpha_1 + \rho \alpha_2 r(1) + \rho (1 - \alpha_1 - \alpha_2) r(2);$$

(2.14)
$$r(2) = \rho \alpha_1 r(1) + \rho \alpha_2 + \rho (1 - \alpha_1 - \alpha_2) r(1);$$

(2.15)
$$r(3) = \rho \alpha_1 r(2) + \rho \alpha_2 r(1) + \rho (1 - \alpha_1 - \alpha_2);$$

(2.16)
$$r(k) = \rho \alpha_1 r(k-2) + \rho \alpha_2 r(k-2) + \rho (1 - \alpha_1 - \alpha_2) r(k-3)$$

for $k \ge 3$. Thus from (2.13) and (2.14)

$$\mathbf{r}(1) = [\rho\alpha_{1} + \rho^{2}(1 - \alpha_{1} - \alpha_{2})\alpha_{2}][1 - \rho\alpha_{2} - \rho^{2}(1 - \alpha_{1} - \alpha_{2})(1 - \alpha_{2})]^{-1};$$

$$r(2) = \rho \alpha_2 + \rho (1 - \alpha_2) r(1);$$

and

$$r(3) = \rho^{2} \alpha_{1} \alpha_{2} + \rho(1 - \alpha_{1} - \alpha_{2}) + \{\rho^{2} \alpha_{1}(1 - \alpha_{2}) + \rho\alpha_{2}\}r(1) .$$

Since equations (2.13)-(2.16) are linear in $\alpha_1' = \rho \alpha_1$, $\alpha_2' = \rho \alpha_2$, and ρ with coefficients involving r(1), r(2) and r(3), they can be solved for α_1' , α_2' and ρ , and thus for α_1 , α_2 and ρ . To facilitate this note that (2.13)-(2.15) can be rewritten as

(2.13')
$$r(1) = \alpha_1^{\prime} \{1 - r(2)\} + \alpha_2^{\prime} [r(1) - r(2)] + \rho r(2);$$

(2.14')
$$r(2) = \alpha'_{2}[1 - r(1)] + \rho r(1);$$

(2.15')
$$r(3) = \alpha_1'[r(2) - 1] + \alpha_2'[r(1) - 1] + \rho$$
.

Hence estimates for ρ , α_1 , and α_2 can be obtained from the usual estimates of the serial correlations r(1), r(2) and r(3). Problems will arise in solving the equations (2.13')-(2.15') if ρ is close to zero, since when $\rho=0$ all the quantities in the equation are identically zero.

The characteristic equation (2.6) becomes

(2.17)
$$0 = 1 - \rho \alpha_1 z - \rho \alpha_2 z^2 - \rho (1 - \alpha_1 - \alpha_2) z^3.$$

Again we require that $\alpha_1 + \alpha_2 < 1$, so that the process does not reduce to the DAR(2) case. To find the roots of (2.17), let

$$a = \frac{1}{3} \left[3 \frac{\alpha_1}{1 - \alpha_1 - \alpha_2} - \left(\frac{\alpha_2}{1 - \alpha_1 - \alpha_2} \right)^2 \right],$$

$$b = \frac{1}{27} \left\{ 2 \left(\frac{\alpha_2}{1 - \alpha_1 - \alpha_2} \right)^2 - 9 \frac{\alpha_1 \alpha_2}{(1 - \alpha_1 - \alpha_2)^2} - 27 \frac{1}{\rho (1 - \alpha_1 - \alpha_2)} \right\}$$

and

$$d = \frac{b^2}{4} + \frac{a^3}{27}$$
.

The value of d can be either positive or negative. If d>0, equation (2.17) has one real positive root and two imaginary roots and the autocorrelation function will have a "damped sine wave" appearance, but the function will eventually decrease geometrically. If d<0, then all the roots of the cubic are real and unequal, one being positive and greater than one, the other two being negative. Hence the autocorrelation function is a mixture of a geometrically decreasing term, and two terms which decrease geometrically in absolute value but oscillate in sign. Table 1 gives values for the first nine correlations for a DAR(3) process with $\rho=0.9$ for various values of α_1 and α_2 .

Example 2.3. The DAR(p) process.

The Yule-Walker equations (2.1)-(2.4) can be reparametrized with $\phi_1 = \rho\alpha_1$, $\phi_2 = \rho\alpha_2$,..., $\phi_p = \rho\alpha_p$, the parameters all lying between 0 and 1, with the constraint that

$$(2.18) 0 \leq \phi_1 + \phi_2 + \cdots + \phi_p < 1.$$

Then it can be seen that the Yule-Walker equations are in exactly the same form as those given for the usual AR(p) process in Box and Jenkins (1970, p. 55). The parameter space for the DAR(p)

TABLE 1

VALUES FOR THE FIRST NINE CORRELATIONS FOR A DAR(3) PROCESS AND α_2 . α**1** WITH $\rho = 0.9$ FOR VARIOUS VALUES OF

α1	^α 2	r(1)	r(2)	r(3)	that α ₁ r(4)	+ α ₂ < r (5)	r(6)	r(7)	r(8)	r(9)
0	0	0	0	6.	0	0	.81	0	0	.729
0	0.5	.58	.71	.71	.58	.64	.58	.55	.55	.51
0	6.0	.40	.85	.41	.72	.41	.62	.40	.54	.38
0.05	6.0	.44	.85	.44	.73	.43	.63	.41	.55	.38
0.1	0.8	.59	.83	.59	.70	.56	.61	.52	.53	.48
0.3	9.0	.75	.81	.71	.70	.64	.61	.58	.54	.51
0.45	0.45	.80	.80	.74	.70	.65	.61	.58	.54	.51
0.5	0	92.	89.	.76	89.	.61	.62	.58	.54	.52
9.0	0.3	.84	.80	.75	69.	. 65	09.	.56	.53	.49
8.0	0.1	.87	.79	.74	89.	.63	.58	.54	.50	.46
 6.0	0	88.	. 79	.73	.67	.62	.57	.52	.48	.44

process is more restricted than that for ϕ_1,\ldots,ϕ_p in the AR(p) case, but the solutions are the same. The parameter ρ does not appear explicitly in this parametrization; however the parametrization is useful in the next section and when negative correlation is considered.

3. ASYMPTOTIC PROPERTIES

3.1. Preliminaries: Aperiodicity and the Doeblin Condition

$$(3.1) P(\underline{i},\underline{k}) \geq (1-\rho)^{p} \pi(k_{1}) \cdots \pi(k_{p}).$$

For C a subset of F let

$$\mu(C) = \sum_{\substack{k \in C}} \pi(k_1) \cdots \pi(k_p) = \sum_{\substack{k \in C}} \mu(k) .$$

The asymptotic theory for DAR(p) processes is based on the following result.

PROPOSITION (3.1). The Markov chain $\{Z_n\}$ is aperiodic and satisfies the Doeblin hypothesis D as generalized by Doob [p. 192, [1953]].

PROOF. Fix a subset C of F such that

$$\delta = \inf_{\mathbf{k} \in \mathbf{C}} \mu(\mathbf{k}) > 0$$
.

Let A be a subset of F with $\mu(A) \leq \mu(C)/2$. Then

$$P(\underline{i}, A^{C} \cap C)$$

$$= \sum_{\underline{k} \in A^{C} \cap C} \frac{P(\underline{i}, \underline{k})}{\mu(\underline{k})} \mu(\underline{k})$$

$$\geq \sum_{\underline{k} \in A^{C} \cap C} (1-\rho)^{p} \mu(\underline{k})$$

$$= (1-\rho)^{p} \mu(A^{C} \cap C)$$

$$\geq (1-\rho)^{p} \mu(C)/2$$

$$\geq (1-\rho)^{p} \delta/2,$$

since $\mu(A) \leq \mu(C)/2$. Hence,

(3.2)
$$P(i,A) = 1 - P(i,A^{C})$$

$$\leq 1 - P(i,A^{C} \cap C)$$

$$\leq 1 - \frac{1}{2} (1 - \rho)^{2} \delta.$$

The result now follows.

We will use proposition (3.1) to show existence of a unique limiting distribution for $\{Z_n\}$, and to show that the stationary DAR(p) process is ϕ -mixing.

3.2. Stationarity Conditions

It follows from Proposition (3.1) that there exists a unique limiting distribution ν for $\{Z_n^{}\}$ and constants $\gamma \geq 0$ and β , $0 \leq \beta \leq 1$ such that

(3.3)
$$|P^{n}(v,A) - v(A)| < \gamma \beta^{n}, \quad n = 1,2,...$$

for all subsets A of F (cf. Doob [1953, p. 221]). A stationary DAR(p) process is obtained by using ν as the initial distribution for $(X_{-p+1}, X_{-p+2}, \ldots, X_0)$. To see that ν has marginal distribution π , let $\{\gamma_n(j), j \in E\}$ denote the marginal distribution of the first component of Z_n and put $\gamma(j) = \lim_{n \to \infty} \gamma_n(j)$. Note that

$$\lim_{n \to \infty} \gamma_n(j) = \sum_{i=1}^{p} \rho \alpha_i \lim_{n \to \infty} \gamma_{n-i}(j) + (1-\rho) \pi(j) .$$

Hence, $\gamma(j) = \rho \gamma(j) + (1-\rho) \pi(j)$ and the marginal distribution of the stationary DAR(p) process is π as asserted in Section 1.

Example 3.1. The DAR(2) process.

Let $\{\textbf{X}_n\}$ be a DAR(2) process with the same parametrization as in Example 2.1. If

(3.4)
$$\gamma(s_1,s_2) = \lim_{n \to \infty} E[\exp\{-s_1 X_{n+1} - s_2 X_{n+2}\}] ,$$
 and
$$\beta(s_1) = \lim_{n \to \infty} E[\exp\{-s_1 X_n\}]$$

then $Y(s_1,s_2)$ satisfies the equation

(3.5)
$$\gamma(s_1, s_2) = (1-\rho) \beta(s_1) \beta(s_2) + \rho\alpha\beta(s_1 + s_2) + \rho(1-\alpha) \gamma(s_1, s_2)$$
.

Solving (3.5) we get

$$\gamma(s_1,s_2) \; = \; \gamma(s_2,s_1) \; = \; (1\; -\; r(1)) \;\; \beta(s_1) \;\; \beta(s_2) \; + r(1) \;\; \beta(s_1+s_2)$$

and inverting the transform results in

(3.6)
$$v(i,j) = \lim_{n \to \infty} P\{X_{n+1} = i, X_{n+2} = j\} = \begin{cases} (1-r(1)) \pi(i) \pi(j), & i \neq j, \\ \\ r(1) \pi(j) + (1-r(1)) \pi(j)^2, & i = j. \end{cases}$$

where $r(1) = \alpha \rho \{1 - \rho (1-\alpha)\}^{-1} = corr(X_n, X_{n+1})$ in the stationary process. Note that $\sum_{j \in E} v(i,j) = \sum_{j \in E} v(j,i) = \pi(i)$, as expected.

In simulating a random variable with distribution ν it is simpler to use the conditional distribution

$$\nu(j|i) = \lim_{n \to \infty} P\{X_{n+2} = j \mid X_{n+1} = i\} = \begin{cases} \{1 - r(1)\} \pi(j), & j \neq i, \\ \\ r(1) + (1-r(1)) \pi(i), & j = i. \end{cases}$$

Note that we can now generate a stationary sequence of random variables X_n , $n=1,2,\ldots$ with any marginal distribution π and

second-order autoregressive (AR(2)) correlation structure. From (3.6) or the conditional distribution it is not hard to show that, with $m = E(Y_n) = E(X_n)$,

(3.7)
$$E[X_n | X_{n-1} = X_{n-1}] = \{1 - r(1)\}m + r(1)X_{n-1},$$

(3.8)
$$E[X_n | X_{n+1} = X_{n+1}] = \{1 - r(1)\} m + r(1) X_{n+1},$$

(3.9)
$$\operatorname{Var}[X_n | X_{n-1} = X_{n-1}] = \{1 - r(1)\} E(X_n^2) + r(1)\{1 - r(1)\} X_{n-1}^2$$

$$- \{1 - r(1)\}^2 m^2 - 2r(1)(1 - r(1)) m X_{n-1}.$$

Directly from (1.1) we have, for example, that

(3.10)
$$E[X_n | X_{n-1} = X_{n-1}; X_{n-2} = X_{n-2}]$$

$$= \rho \alpha X_{n-1} + \rho (1-\alpha) X_{n-2} + (1-\rho) m;$$

(3.11)
$$\operatorname{Var}[X_{n} | X_{n-1} = X_{n-1}; X_{n-2} = X_{n-2}]$$

$$= (1-\rho) \operatorname{Var}(Y_{n}) + \rho\alpha(1-\rho\alpha) X_{n-1}^{2} \qquad (1-\rho(1-\alpha)) X_{n}^{2}$$

$$+ (1-\rho) \rho E[Y_{n}^{2}] + 2\rho^{2}\alpha(1-\alpha) X_{n-1} X_{n}$$

$$+ 2\rho(1-\rho) m[\alpha X_{n-1} + (1-\alpha) X_{n}] .$$

The regressions (3.7) and (3.8) might suggest that the DAR(2) process is time-reversible, but this is not so. To see this, note that

$$\begin{array}{l} \gamma(s_{1},s_{2},s_{3}) &= \lim\limits_{n\to\infty} \ \mathrm{E}[\exp\{-s_{1}^{X}_{n+1} - s_{2}^{X}_{n+2} - s_{3}^{X}_{n+3}\}] \\ \\ &= \rho\alpha\gamma(s_{1},s_{2}+s_{3}) + \rho(1-\alpha) \ \gamma(s_{1}+s_{3},s_{2}) \\ \\ &+ (1-\rho) \ \beta(s_{3}) \ \gamma(s_{1},s_{2}) \end{array}.$$

Since $\gamma(s_1, s_2, s_3) \neq \gamma(s_3, s_2, s_1)$ it follows that the stationary DAR(2) process is not time reversible.

Example 3.2. The DAR(3) process.

Let $\{x_n^{}\}$ be a DAR(3) process with the same parametrization as in Example (2.2). As before let

$$\gamma(s_1,s_2) = \lim_{n \to \infty} E[exp\{-s_1^{X}_{n+1} - s_2^{X}_{n+2}\}]$$
,

and let

$$\beta(s_1) = \lim_{n \to \infty} E[\exp\{-s_1^{X}_n\}];$$

$$\delta(s_1,s_2) = \lim_{n \to \infty} E[exp\{-s_1^{X}_{n-1} - s_2^{X}_{n+1}\}];$$

and

$$\Gamma(s_1, s_2, s_3) = \lim_{n \to \infty} E[\exp\{-\{s_1^{X}_{n+1} - s_2^{X}_{n+2} - s_3^{X}_{n+3}\}].$$

Put $\phi_1 = \rho \alpha_1$, $\phi_2 = \rho \alpha_2$, and $\phi_3 = \rho (1 - \alpha_1 - \alpha_2)$. Then, from the definition (1.1), and accounting for the possibility of X_{n+3} taking values X_{n+2} , X_{n+1} , X_n or Y_{n+3} , we have

(3.12)
$$\Gamma(s_{1}, s_{2}, s_{3})$$

$$= (1-\rho) \beta(s_{3}) \gamma(s_{1}, s_{2}) + \phi_{1}\gamma(s_{1}, s_{2}+s_{3})$$

$$+ \phi_{2}\gamma(s_{1}+s_{3}, s_{2}) + \phi_{3} \Gamma(s_{3}, s_{1}, s_{2}) .$$

Also

$$\begin{array}{lll} (3.13) & \gamma(s_1,s_2) \\ \\ &= (1-\rho) \ \beta(s_1) \ \beta(s_2) \ + \ \phi_1\beta(s_1+s_2) \ + \ \phi_2\gamma(s_2,s_1) \ + \ \phi_3\delta(s_1,s_2) \end{array}$$

and

$$\begin{array}{lll} & \delta \left(\mathbf{s_{1}, s_{2}} \right) \\ \\ & = (1 - \rho) \ \beta \left(\mathbf{s_{1}} \right) \ \beta \left(\mathbf{s_{2}} \right) \ + \ \phi_{1} \gamma \left(\mathbf{s_{1}, s_{2}} \right) \ + \ \phi_{2} \gamma \left(\mathbf{s_{1} + s_{2}} \right) \ + \ \phi_{3} \gamma \left(\mathbf{s_{2}, s_{1}} \right). \end{array}$$

Solving (3.13) and (3.14) simultaneously for $\gamma(s_1,s_2)$ and $\delta(s_1,s_2)$, it follows that

$$\gamma(s_1,s_2) = (1 - r(1)) \beta(s_1) \beta(s_2) + r(1) \beta(s_1 + s_2)$$
,

where $r(1) = corr(X_n, X_{n+1})$ is given in Example (2.2). Hence, inverting the transform,

(3.15)
$$v(i,j) = \begin{cases} (1-r(1)) \pi(i) \pi(j), & \text{if } i \neq j, \\ \\ r(1) \pi(j), & \text{if } i = j. \end{cases}$$

The expression (3.12) can now be used to obtain an expression for $\Gamma(s_3,s_1,s_2)$, and substituting back into (3.12) we get

Note that $\Gamma(s_1, s_2, s_3) \neq \Gamma(s_3, s_2, s_1)$ in general and hence the stationary DAR(3) process is not time reversible.

3.3. Mixing Results

The following result follows directly from (3.3).

<u>PROPOSITION 3.2.</u> The stationary DAR(p) process is ϕ -mixing in the sense of Billingsley [1968] with $\phi(k) = \gamma \beta^k$.

This last result implies that all of the asymptotic results in Jacobs and Lewis [1978b] are true for the DAR(p) process modulo constants. In particular the usual estimates of covariances, percentiles and quantiles converge almost surely to the true covariances, percentiles and quantiles and the estimates are asymptotically normal. Further, a chi-square goodness of fit test can be obtained for the marginal distribution of the stationary DAR(p) process. For further details see Jacobs and Lewis [1978b].

4. EXTENSIONS

One possible drawback of the DAR(p) process as defined in (1.1) is that all of the correlations are positive. We will now discuss a variation of the DAR(p) process which allows negative correlation for the case in which the marginal distribution π is symmetric about zero. The case of nonsymmetric π is more difficult and will be considered elsewhere.

Let $\{v_n^{}\}$, $\{\mathtt{A}_n^{}\}$, and $\{\mathtt{Y}_n^{}\}$ be as in Section 1 and suppose that π is symmetric about zero. Let

(4.1)
$$X_n = V_n a_{A_n} X_{n-A_n} + (1 - V_n) Y_n$$
,

where $\{a_k; k = 1,...,p\}$ is a fixed sequence of numbers that are either -1 or 1.

Put $Z_n = (X_n, X_{n+1}, \dots, X_{n+p-1})$. Then $\{Z_n\}$ is a Markov chain with a transition matrix P which satisfies (3.1). Thus Propositions (3.1) and (3.2) still hold, as well as (3.3).

Let $\{X_n^{}\}$ be a stationary process defined by the modified DAR(p) scheme of (4.1) and put $r(k) = corr(X_n^{}, X_{n+k}^{})$. The Yule-Walker equations for this process are

(4.2)
$$r(1) = \phi_1 + \phi_2 r(1) + \cdots + \phi_p r(p-1);$$

(4.3)
$$r(2) = \phi_1 r(1) + \phi_2 + \cdots + \phi_p r(p-2);$$

$$\vdots \qquad \vdots \qquad \vdots \qquad \vdots$$
(4.4)
$$r(p) = \phi_1 r(p-1) + \phi_2 r(p-2) + \cdots + \phi_p r(0);$$

(4.4)
$$r(p) = \phi_1 r(p-1) + \phi_2 r(p-2) + \cdots + \phi_p r(0);$$

(4.5)
$$r(p+k) = \phi_1 r(p+k-1) + \phi_2 r(p+k-2) + \cdots + \phi_p r(k)$$
;

for $k \ge 1$ where $\phi_k = a_k \rho \alpha_k$, k = 1, ..., p.

Equations (4.2)-(4.4) have the same form as the Yule-Walker equation on page 55 of Box and Jenkins (1970) but the set of possible values of ϕ_i , i = 1, ..., p is more restrictive. In particular we always have that

$$|\phi_1| + |\phi_2| + \cdots + |\phi_p| < 1$$

Example 4.1. The modified DAR(1) process.

Let $A_n \equiv 1$ and $a_1 = -1$. Then

$$X_{n+1} = -V_{n+1}X_n + (1 - V_{n+1})Y_{n+1}$$

and $r(k) = (-1)^k \rho^k$. If $a_1 = 1$ the process is the same as that in Section 1. When $a_1 = -1$ the modified DAR(1) process is Markovian with transition matrix

$$P(i,j) = P\{X_{n+1} = j | X_n = i\} = \begin{cases} (1-\rho) & \pi(j), & -i \neq j, \\ \\ \rho + (1-\rho) & \pi(j), & -i = j. \end{cases}$$

Example 4.2. The modified DAR(2) process.

Let $P\{A_n = 1\} = 1 - P\{A_n = 2\} = \alpha$, $0 \le \alpha \le 1$. Solving equations (4.2)-(4.4) we obtain

$$r(1) = a_1^{\rho\alpha} [1 - a_2^{\rho} (1-\alpha)]^{-1},$$

$$r(2) = \rho [a_2^{\rho} (1-\alpha) - \rho (1 - 2\alpha)] [1 - a_2^{\rho} (1-\alpha)]^{-1},$$

and

$$r(3) = \rho^2 \alpha [2a_1a_2(1-\alpha) - a_1\rho(1-2\alpha)] [1 - a_2\rho(1-\alpha)]^{-1}$$
.

Note that if $a_1 = 1$ (respectively $a_1 = -1$), the r(1) > 0, (respectively r(1) < 0) and r(2) can be either positive or negative. The analogue of the characteristic equation (2.6) is

(4.6)
$$1 - a_1 \rho \alpha z - a_2 \rho (1-\alpha) z^2 = 0.$$

If $a_2 = 1$, then (4.6) has two real roots; if $a_2 = -1$, then (4.6) may have imaginary roots. Thus the autocorrelation function can either have a "damped sine wave" appearance or be a mixture of terms which decrease geometrically in absolute value but may oscillate in sign.

For the modified DAR(2) process the range of values which the parameters ϕ_1 and ϕ_2 can take is not as broad as for the AR2 case, for which we have (Box and Jenkins [1970, p. 58])

which allows ϕ_1 to range between -2 and +2. By contrast for the modified DAR(2) (for symmetric distributions) we have

Thus we have a diamond of possible values in the (ϕ_1,ϕ_2) plane, with apexes (0,1), (1,0), (0,-1), (-1,0). With this restriction the behavior of the serial correlations for the modified DAR(2) process is the same as the behavior for the AR(2) process which is pictured in the figure in Box and Jenkins (1970).

A limiting stationary distribution can be derived for this modified DAR(2) process, as was done for the ordinary DAR(2) process in Example 3.1.

REFERENCES

- Billingsley, P. (1968). <u>Convergence of Probability Measures</u>. New York: Wiley.
- Box, G.E.P. and Jenkins, G.M. (1970). <u>Time Series Analysis</u>. San Francisco: Holden-Day.
- Doob, J. L. (1953). Stochastic Processes. New York: Wiley.
- Jacobs, P.A. and Lewis, P.A.W. (1978a). Discrete Time Series Generated by Mixtures.I: Correlational and Runs Properties. J.R. Statist. Soc. B, 40, 94-105.
- Jacobs, P.A. and Lewis, P.A.W. (1978b). Discrete Time Series Generated by Mixtures. II: Asymptotic Properties.

 J.R. Statist. Soc. B, 40, to appear.

DISTRIBUTION LIST

			NO.	OF	COPIES
	STATISTICS AND FRCBABILITY CFFICE OF NAVAL RESEARCH CODE 436	PROGRAM			
	ARLING TON	22217		1	
	CFFICE OF NAVAL RESEARCH NEW YORK AREA CFFICE 715 BROACWAY - 5TH FLOOR ATTN: DR. RJEER GRAFTON				
	NEW YORK, NY	10003		1	
	DIRECTOR OFFICE OF NAVAL RESEARCH B	RANCH OFF			
	ATTN: DEPUTY AND CHIEF SC CHICAGO, IL	1 ENT 1ST 60605		1	
,	LIBRARY NAVAL OCEAN SYSTEMS CENTER SAN DIEGO				
	CA	92152		1	
	NAVY LIBRARY NATIONAL SPACE TECHNOLOGY I ATTN: NAVY LIBRARIAN	LAB			
	EAY ST. LEUIS	39522		1	
	NAVAL ELECTRONIC SYSTEMS CO	GNAMMO			
	NATICNAL CENTER NO. 1 ARLINGTON VA	20360		1	
	CIRECTOR NAVAL REASARCH LAN	EURATORY			
	CODE 2029 WASHINGTON, C.C.	20375		1	
	DEFENSE COCUMENTATION CENTS CAMERON STATION	ER		2	
	ALEXANDRIA Vifginia 22314				
	TECHNICAL INFORMATION DIVIS NAVAL RESEARCH LABORATORY	SION		1	
	WASHINGTON. C. C.	20375			

C F	FI	C	E	D	F	N	4	V 1	١	0	R	ES	EA	R	C	+							NO.	0	F	C	OPI	E	S
76 SA	000	M F	RA	KA	Č	S	Š	TR C	E	EA	ŤL	_ I F	: 01	315	11	Δ		9	41	02				1					
TE NA I N	VA	1 L	0	R		A	N	CE	:	S	T					3 6	4	0						1					
NA PH DI FH	l L VI	S	36	L	FH	I	A Ci	41	ΙI	С	ΔΙ		LI	8	R	ΔR	Y			91	12			J					
BU DE TE	C٢	IN.	IC	4	L	L	I	ER	Δ	₹'	Υ		SC AV											1	L,				
LI NA MC	VA	L	P	0	ST	G	R.	5 C	L	Δ.	T						4	٥							ı				
PR DE LN CH NC	FA I V	REF	MS	E	TY TY	- (C	=	M	A	T	4 E	MA	T			I		A 282	223	3			1					
P DE ST	FA	FC	IM IR	E!	U	1	IV	: / E	255	5	I -	IY	ST	1			3(05	5					1					
F NE	LE k	HA	N Vi	I I	15 1	R	5 1	T	Υ				BEST	I	С	S								1					
PRI IN IN IN IN IN	I O	NE	S		EAL	E	3 2	N		I	35	T	RI	A										1					
PRE UN CHO	FAVPR	RTER	CMS	ENTAR	RTYLL		BEE	A	K!	CI	AT	I H	ST	IA	C	S	11	4 4	8					,					

PROF. R. E. BECHGFER DEFARTMENT OF OPERATIONS RESEARCH CORNELL UNIVERSITY ITHACA NEW YORK 1485C	1
FRCF. N. J. BERSHAD SCHOOL OF ENGINEERING UNIVERSITY OF CALIFORNIA IRVINE . CALIFORNIA \$2664	1
P. J. BICKEL CEPARTMENT OF STATISTICS UNIVERSITY OF CALIFORNIA BEFKELEY . CALIFORNIA	1
94720	
PROF. H. W. BLCCK DEFARTMENT OF MATHEMATICS UNIVERSITY OF FITTSBURGH PITTSBURGH PA	1
15263	
FROF. JOSEPH BLUM DE FT. OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE THE AMERICAN UNIVERSITY MASHINGTON DC 20016	1
PFOF. R. A. BRADLEY	1
DEPARTMENT OF STATISTICS FLORIDA STATE UNIVERSITY	
TALLAHASSEE , FLORIDA 323C6	
PROF. R. E. BARLOW CFERATIONS RESEARCH CENTER CCLLEGE OF ENGINEERING LNIVERSITY OF CALIFORNIA BEFKLEY CALIFORNIA 54720	1
MR. C. N BENNETT NAVAL COASTAL SYSTEMS LABORATORY CODE P761 PANAMA CITY, FLCRIDA 32401	1
PROF. U. N. BEAT CCMPUTER SCIENCE / OPERATIONS RESEARCE CENTER SOUTHERN METHODIST UNIVERSITY CALLAS TEXAS 75275	1
PRCF. W. R. BLISCHKE DEPT. OF QUANTITATIVE BUSINESS ANALYSIS UNIVERSITY OF SOUTHERN CALIFORNIA LOS ANGELES, CALIFORNIA 50007	1

	NO. OF COPIES
DR. DERRILL J. BORDELON NAVAL UNCERWATER SYSTEMS CENTER COCE 21 NEWPORT	•
RI C2 84 0	
J. E. BOYER JR DEPT. OF STATISTICS SCUTHERN METHOLIST UNIVERSITY DALLAS TX	1
75275	
DR. J. CHANERA U. S. ARMY PESEARCH P. G. BCX 12211 RESEARCH TRIANGLE PARK, NCFTH CAROLINA 27706	
PROF. L. CHERNOFF DEPT. OF MATHEMATICS MASS INSTITUTE OF TECHNOLOGY CAMBRIDGE. MASSACHUSETTS 02139	1
PROF. C. DERMAN DEFARTMENT OF CIVIL ENGINEERING AND ENGINEERING MECHANICS COLUMBIA UNIVERSITY NEW YORK NEW YORK 10027	1
FRCF. R. L. CISNEY VIRGINIA PCLYTECHNIC INSTITUTE AND STATE UNIVERSITY DEPT. CF INDLSTRIAL ENGINEERING AND CPERATIONS RESEARCH BLACKSBURG, V/ 24061	1
PR. J. DOWLING DEFENSE LOGISTICS STUDIES INFORMATION EXCHANGE ARMY LOGISTICS MANAGEMENT CENTER FORT LEE VIRGINIA 20390	1
PRCF J. D. ESARY CEPT. CF OFERATIONS RESEARCH AND ADMINISTRATIVE SCIENCES NAVAL POSTGRADUATE SCHOOL MONTEREY CALIFORNIA 93940	1
DR. M. J. FISCHER DEFENSE COMMUNICATIONS AGENCY 1860 WIEHLE AVENUE RESTON VIRGINIA 22070	1

PROF. C. P. GAVER DEFT. OF OPERATIONS RESEARCH NAVAL FOSTGRADUATE SCHOOL MONTEREY CA	1
93940	
MR. GENE F. GLEISSNER APPLIED MATHEMATICS LAEDRATORY CAVID TAYLOR NAVAL SHIP RESEARCH AND DEVELOPMENT CENTER BETHESCA MD 20084	1
PROF. S. S. GUPTA DEPARTMENT OF STATISTICS FURDUE UNIVERSITY LAFAYETTE INDIANA 47907	1
PROF. D. L. HANSON DEFT OF MATH. SCIENCES STATE UNIVERSITY OF NEW YORK, EINGHAMTON BINGHAMTON NY 13901	1
PROF. F. J. HARRIS DEFT. OF ELECTRICAL ENGINEERING SAN CIEGO STATE UNIVERSITY SAN DIEGO CA \$2182	1
FRCF. L. H. HERBACH DEFT. CF CFERATICAS RESEARCH AND SYSTEMS ANALYSIS POLYTECHNIC INSTITUTE CF NEW YORK BROOKLYN NY 11201	1
PRCF. M. J. HINICH DEPARTMENT OF ECONOMICS VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY BLACKSBURG, VIRGINIA 24C61	1
PROF. W. M. HIRSCH INSTITUTE OF MATHEMATICAL SCIENCES NEW YORK UNIVERSITY NEW YORK NEW YORK 10453	1
PROF. C. L. ICLEHART DEFARTMENT OF OPERATIONS RESEARCH STANFORD UNIVERSITY STANFORD, CALIFORNIA 94350	1

PRCF. J. B. KACANE DEPARTMENT CF STATISTICS CARNEGIE-MELLCN FITTSBURGH, PENNSYLVANIA 15213	1
DR. RICHARE LAU DIRECTOR CFFICE OF NAVAL RESEARCH BRANCH OF 1020 EAST GREEN STREET FASADENA CA 91101	. 1 F
DR. A. R. LAUFER DIRECTOR OFFICE OF NAVAL RESEARCH ERANCH OF 1030 EAST GREEN STREET PASADENA CA 91101	1 F
PROF. M. LEACEETTER DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CAPOLINA CHAPEL HILL NORTH CAROLINA 27514	1
CR. J. S. LEE J. S. LEE ASSCCIATES, INC. 2001 JEFFERSON DAVIS HIGHWAY SUITE EC2 ARLINGTON VA 22202	1
PROF. L. C. LEE DEPARTMENT OF STATISTICS VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY BLACKSBURG VA 24061	1
PROF. R. S. LEVENWORTH DEPT. OF INCUSTRIAL AND SYSTEMS ENGINEERING UNIVERSITY OF FLORIDA GAINSVILLE FLORIDA 22211	
PRCF G. LIEBERMAN STANFORD INIVERSITY DEPARTMENT OF CPERATIONS RESEARCH STANFORD CALIFORNIA 94305	. 1

DR. JAMES F. MAAR NATIGNAL SECURITY AGENCY FORT MEADE, MARYLAND 20755	1
PRCF. R. W. MACSEN DEFARTMENT OF STATISTICS UNIVERSITY OF MISSOURI COLUMBIA MC 65201	1
DR. N. R. MANN SCIENCE CENTER RCCKWELL INTERNATIONAL CORPORATIC F.C. BCX 1085 THOUSAND CAKS, CALIFORNIA 91360	1 N
DR. W. H. MARLOW PREGRAM IN LOGISTICS THE GEERGE FASHINGTON UNIVERSITY 707 22ND STREET, N. W. WASHINGTON, E. C. 20037	1
PROF. E. MASRY DEFT. APPLIEC PHYSICS AND INFORMATION SERVICE UNIVERSITY OF CALIFORNIA LA JOLLA CALIFORNIA \$2093	1
DF. BRUCE J. MCCONALD SCIENTIFIC DIRECTOR SCIENTIFIC LIAISON GROUP CFFICE OF NAVAL RESEARCH AMERICAN EMPASSY - TOKYO AFC SAN FRANCISCO 96503	1
FRCF. J. A. MUCKSTADT CEPT. CF CFERATIONS RESEARCH CCRNELL UNIVERSITY ITHACA, NEW YORK 19850	1
DF. JANET M. MYFRE THE INSTITUTE OF DECISION SCIENCE FOR BUSINESS AND PUBLIC FCLICY CLAREMONT PEN'S COLLEGE CLAREMONT CA 91711	1
PR. H. NISSELSON BLREAU OF THE CENSUS PCCM 2C25 FROERAL FUILCING 3 WASHINGTON . C. C. 2C33	1

MISS B. S. ORLEANS NAVAL SEA SYSTEMS COMMAND (SEA 03F) RM 10508 ARLINGTON VIRGINIA 20360	1
PROF. C. B OWEN CEPARTMENT OF STATISTICS SCUTHERN METHODIST UNIVERSITY DALLAS TEXAS 75222	1
DR. A. PETRASOVITS ROOM 2078 , FGCD AND DRUG ELDG. TENNSY'S PASTURE OTTOWA , CNTARIC K1A-GL2 , CANADA	1
PROF. S. L. PHCENIX SIBLEY SCHOOL OF MECHANICAL AND AEROSPACE ENGINEERING CCFNELL UNIVERSITY ITHACA NY 14850	1
DR. A. L. POWELL CIRECTOR CFFICE OF NAVAL RESEARCH BRANCH OFF 495 SUMMER STREET BOSTON VA C2213	1
MR. F. R. PRIORI CCCE 224 CPERATIONSL TEST AND ONRS EVALUATION FORCE (OPTEVFOR) NORFOLK , VIFGINIA 20360	1
PROF. W. L. FURI DEFT. CF MATHEMATICS F.C. BOX F INCIANA UNIVERSITY FOUNDATION BLOOMING TON IN 47401	1
PROF. H RCBBINS DEFARTMENT OF MATHEMATICS COLUMBIA UNIVERSITY NEW YORK, NEW YORK, NEW YORK,	1

FROF. M ROSENBLATT DEPARTMENT OF MATHEMATICS UNIVERSITY OF CALIFORNIA SAN DIEGO LA JOLLA CALIFORNIA \$2093	1
FROF. S. M. ROSS COLLEGE OF ENGINEERING UNIVERSITY OF CALIFORNIA BEFKELEY CA 54720	1
PRCF. I RUBIN SCHOOL OF ENGINEERING AND APPLIEC SCIENCE UNIVERSITY OF CALIFCRNIA LCS ANGELES . CALIFCRNIA 90024	1
PROF. I. R. SAVAGE DEFARTMENT OF STATISTICS YALE UNIVERSITY NEW HAVEN. CONNECTICUT (652)	1
PROF. L. L., JR. SCHARF DEFARTMENT OF ELECRICAL ENGINEERING COLORACO STATE UNIVERSITY FT. COLLINS, COLDRADO 80521	1
FRCF. R. SERFLING DEPARTMENT OF STATISTICS FLORIDA STATE UNIVERSITY 14LLAHASSEE FLORIDA 32306	1
PROF. N. R. SCHUCANY DEPARTMENT OF STATISTICS SCUTHERN METHOLIST UNIVERSITY DALLAS, TEXAS 75222	1
FRCF. C. O. SIEGMUND DEFT. CF STATISTICS STANFORD UNIVERSITY STANFORC CA 94305	. 1
FRCF. M. L. SHOOMAN DEFT. CF ELECTRICAL ENCINEERING FOLYTECHNIC INSTITUTE OF NEW YORK BROOKLYN, NEW YORK 11201	
FRCF. N. SINGPURWALLA DEPT. CF CFERATIONS RESEARCH THE GEORGE WASHINGTON UNIVERSITY 707 22ND ST. N. W. WASHINGTON, C. C. 20052	1

DR. A. L. SLAFKOSKY SCIENTIFIC ADVISOR COMMANCANT OF THE MARINE CORPS WASHINGTON . C. C. 20380	1
MR. CHARLES S. SMITH DASD (IGL), FENT AGON AASHINGT CN DC 20301	1
20301	
DR. D. E. SMITH DESMATICS INC. P.C. BCX 618 STATE COLLEGE PENNSYLVANIA 16801	1
PRCF. W. L. SMITH DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CARCLINA CHAPEL HILL NOFTH CAROLINA 27514	1
FRCF. H SOLOMON DEFARTMENT OF STATISTICS STANFORD UNIVERSITY STANFORC , CALIFERNIA 94305	1
MR. GLENN F. STAHLY NATIONAL SECURITY AGENCY FORT MEADE MARYLAND 20755	1
MR. CAVIC A. SWICK ACVANCED PROJECTS GROUP CODE 8102 NAVAL FESEARCH LAB. NASHINGTON OC 20375	1
NO WENDELL C SYVES	
PR. WENDELL G. SYKES ARTHUR D. LITTLE, INC. ACCRN PARK CAMBRICGE MA	1
C2140	
FROF. J. R. THOMPSON DEPARTMENT OF MATHEMATICAL SCIENCE RICE UNIVERSITY HOUSTON, TEXAS 77001	1

	NO.	OF	COPIES
FRCF. h. A. THOMPSON DEPARTMENT OF STATISTICS UNIVERSITY OF MISSOURI COLUMBIA, MISSOURI 65201		1	
FRCF. F. A. TILLMAN DEPT. DF INCUSTRIAL ENGINEERING KANSAS STATE UNIVERSITY MANHATTAN		1	
KS 66506			
FRCF J. W. TUKEY DEFAFTMENT OF STATISTICS PRINCETON UNIVERSITY		1	
PRINCETON . N. J. C854C			
FROF. A. F. VEINOTT CEFARTMENT OF OPERATIONS RESEARCH STANFORD UNIVERSTITY STANFORD CALIFORNIA 54305		1	
CANTEL H. WAGNER STATION SCLARE CHE FACLI . PENNSYLVANIA 19301		1	
PROF. GRACE WAHBA DEPT. OF STATISTICS UNIVERSITY OF WISCONSIN MADISON L 53706		1	
FRCF. K. T. WALLENIUS DEPARTMENT OF MATHEMATICAL SCIENCES CLEMSON UNIVERSITY CLEMSON, SOLTH CAROLINA 29631		1	
PROF. G. S. MATSON CEPARTMENT OF STATISTICS FRINCETON, N. J. 08540		1	
FRCF. BERNAFC WICROW STANFORD ELECTRONICS LAB STANFORD UNIVERSITY STANFORD CA 543 05		1	
FRCF. G. E. WHITEHOUSE DEPT. OF INDUSTRIAL ENGINEERING LEHIGH UNIVERSITY BETHLEHEP FA		1	
• • •			

PROF. S. ZACKS CEFT. OF MATHEMATICS AND STATISTICS CASE WESTERN FESERVE UNIVERSITY CLEVELAND., CHIC 44106	1
PRCF. M. ZIA-FASSAN DEFARTMENT OF INDUSTRIAL AND SYSTEMS ENGINEERING ILLINCIS INSTITUTE OF TECHNOLOGY CFICAGO IL	1
CR. DAVID BRILLINGER STATISTICS DEPT. UNIVERSITY OF CALIFORNIA BEAKELEY, CALIFORNIA 54720	1
CR. CHIN CHEN CCMMONWE ALTH ASSOCIATES, INC. 209 E. WASHINGTON JACKSON, MI 49201	1
FRCFESSOR PETER CHEN ALFRED P. SLCAN SHOOOL OF MGNT MASS INSTITUTE OF TOHNOLOGY 50 MEMORIAL DRIVE CAMBRICGE; MASSACHUSETTS 02139	1
DR. E. CINLAR O.R. DEPT. ACRTHMESTERN UNIVERSITY EVANSTON , ILLINOIS 60201	1
DR. GUY C CORYNEN L. 156 UNIV OF CALIFORNIA LAWRENCE LIVERMORE LABORATORY F. D. EOX 808 LIVE FMCRE, CALIFORNIA 94550	1
DR. MICHAEL A. CRANS CENTROL ANALYSIS CORPORATION EOC WELCH READ FALO ALTO, CALIFORNIA 94304	1
CR. D. F. CALEY STATISTICS DEPT. (I A S) AUSTRALIAN NATIONAL UNIVERSITY CANBERRA, A. C. T. 2606 AUSTRALIA	. 1

HSU DER-ANN SCH OF ENGF./APPLIED SCI. DEFT. OF CIVIL & GEOLOGICAL ENGR. THE ENGINEERING QUADRANGLE FRINCEION NEW JERSEY C8543	1
ASSISTANT CEAN CAV ELDREDGE SCHOOL OF EUSINESS ADMINISTRATION UNIVERSITY OF EVANSVILLE EVANSVILLE 1 NDIANA 47732	1
CR. G. W. FISHMAN CIFRICULUM IN O. R. UNIVERSITY OF NORTH CARCLINA PHILLIPS ANNEX, CHAPPEL HILL NORTH CAROLINA 27514	1
CR. A. V. FOLCEN DEPARTNENT OF PHYSIOLOGY THE UNIVERSITY LEEDS LS2 SJT ENGLANC	1
DR. ELGENE C. FOMER MANAGEMENT SCI. & ENGINEERING DEPT. C. W. PCST COLLEGE CF LONG ISLAND UNIVERSITY GREENVALE, NEW YCRK 11548	1
DR. D. VERE JONES DEPT. OF MATHEMATICS VICTORIA UNIVERSITY OF WELLINGTON , P. D. BCX 196 MELLINGTON , NEW ZEALAND	1
J. M. LIITTSCHHAGER SYSTEMS CIVISION CCLLEGE OF ENGINEERING THE UNIVERSITY OF IOWA IDHA CITY , ICHA 52242	1
Y. W. CE KHAADSTENIET DE FATRMENT OF EIULOGY FREE UNIVERSITY CE BOELELAAN 1087 AMSTERCAM-EUITENVELDER THE NETHERLANDS	1
JACK P. LANCOLT DEFENCE & CIVIL INST. OF ENVIR. MED 1133 STEPART AVE. WEST PO BOX 2000 DOWNSVIEW ONTARIO. CANADA	1

MF. FRANK MCNOLTY 1142 FCME AVE. SUNNYVALE, CA 94087	1
CR. P. A. F. VCRAN STATISTICS CEPT. (I A S) AUSTRALIAN NATIONAL UNIVERSITY CANBERRA, A. C. T 2606 AUSTRALIA	1
G. W. MORRISON DAK RIDGE NATIONAL LABORATORY CAK RIDGE, TENNESSE 37830	1
MR. BRADLEY NICKEY LSCA. FOREST SERVICE P. 0. BOX 5007 FIVERSIDE CALIFORNIA 92507	1
CR. DAVID CAKES DEFT. OF STATISTICS HARVARE UNIVERSITY SCIENCE CENTER, 1 OXFORD ST. COMBRIDGE, MA. 02132	1
Professor Emanuel Parzen, Director Institute of Statistics Teaxs A&M University College Station, TX 77843	1
MR ART PETERSON DEFT. OF EIGSTATISTICS SC-32 UNOVERSITY OF WASHINGTON SEATTLE, WASHINGTON 58195	1
CAVID M. RCBINSON CONTROL ANALYSIS CCRPORATION 800 WELCH ROAD PALD ALTO CALIFORNIA \$4304	1
C. H. SAUER (21-129) THOMAS J. WATSCN RESEARCH CENTER F.C. BCX 218 YCRKTOWN HEIGHTS, NEW YCRK 10598	1

MR. C. A. STEELE JR F.C. BCX 45 MACNCLIA, MA C1930	1
PRCF P. SULLO SCHOOL OF MANAGEMENT FENSSELAER POLYTECHNIC INSTITUTE TROY, N.Y. 12181	1
DR. FOBERT TRAPPL INSTITUT FUR ALLGEMEINE UND VERGLEICHENDE PHYSIOLOGIE DER UNIVERSITAT WIEN SCHWARZSPANIERSTRASSE 17 A-1090 WIEN GSTERREICH (ALSTRIA)	1
PALL A. HILLIS PCLY TECHNICA 2824 W. GECRGE MASON FALLS CHURH, VA. 22042	1
TUNCEL M. YEGULALP KRLMB SCHCCL CF MINES 918 S.h. MLDD CCLUMBIA UNIVERSITY 520 h. 12CTH ST. NEW YROK, N.Y. 10027	1
SHIEGKI YOKOYAMA RESEARCH CENTER OF APPLIED INFORMATION SCIENCES TOHOKU UNIVERSITY KATAHIRA-CHO 2-1, SENDAI 980 JAPAN	1
Dean of Research Code 012 Naval Postgraduate School Monterey, Ca. 93940	1
Library (Code 55) Naval Postgraduate School Monterey, CA 93940	1

Code 55Lw	50
Naval Postgraduate School Monterey, CA 93940	
Monterey, CA 93940	
Professor P. A. Jacobs Code 55Jc	50
Naval Postgraduate School	
Monterey, CA 93940	
R. J. Stampfel Code 55	1
Naval Postgraduate School	
Monterey, CA 93940	

